

Claims:-

1. A method of producing a desired protein, polypeptide or peptide, comprising the step of:
- 5 culturing a mammalian host cell in culture medium, wherein said host cell includes:
- (i) at least one introduced DNA sequence encoding the desired protein, polypeptide or peptide expressibly linked to a first inducible promoter sequence,
- 10 (ii) at least one introduced DNA sequence encoding a protein, polypeptide and/or peptide factor(s) required for growth of the host cell in said culture medium expressibly linked to a promoter sequence, the expression of which is regulated by a repressor binding region; and
- (iii) at least one introduced DNA sequence encoding a repressor molecule which binds to the repressor binding region, expressibly linked to
- 15 a second inducible promoter sequence.
- wherein the first and second inducible promoter sequence(s) may be the same or different.
- 20 2. A method according to claim 1, wherein the first and second, inducible promoter sequence(s) are the same, and said culturing step comprises:
- a first stage of culturing said host cell to a desired cell confluence, and a second stage of culturing said host cell in the presence of an inducer of
- 25 said first and second inducible promoter sequence(s).
3. A method according to claim 1 or 2, wherein the said repressor binding region is a *lac* operator sequence, and said at least one DNA sequence encoding a repressor molecule encodes a *lac* repressor.

4. A method according to any one of claims 1 to 3, wherein the first and second, inducible promoter sequence(s) is/are selected from the group consisting of the human metallothionein IIA promoter and the modified human metallothionein IIA promoters, M(1)2 and M(2)6.

5. A method according to claim 4, wherein the host cell further includes and expresses a DNA sequence encoding a metallothionein.

6. A method according to any one of the preceding claims, additionally comprising the step of recovering the desired protein, polypeptide or peptide.

Sub All 7. A method for the regulated growth of a mammalian host cell in a culture medium, comprising the step of:-

15 culturing said mammalian host cell in said culture medium, wherein said host cell includes:

(i) at least one introduced DNA sequence encoding a protein, polypeptide and/or peptide factor(s) required for growth of the host cell in said culture medium expressibly linked to a promoter sequence, the expression of which is regulated by a repressor binding region; and

20 (ii) at least one DNA sequence encoding a repressor molecule which binds to the repressor binding region, expressibly linked to an inducible promoter sequence.

25 8. A method according to claim 7, wherein the said repressor binding region is a *lac* operator sequence, and said at least one DNA sequence encoding a repressor molecule encodes a *lac* repressor.

9. A method according to claims 7 or 8, wherein the inducible promoter sequence(s) is/are selected from the group consisting of the human

09564338-092801

10. A method according to claim 9, wherein the host cell further
5 includes and expresses a DNA sequence encoding a metallothionein.

12. A method according to claim 11, wherein the DNA sequence(s)
encoding the protein, polypeptide and/or peptide growth factor(s) encodes
15 insulin or an insulin-like growth factor.

20

Sub A12

25

16. A method according to any one of the preceding claims, wherein the mammalian host cell is a CHO-K1 cell.

17. A host cell including:

(i) at least one introduced DNA sequence encoding a desired protein, polypeptide or peptide expressibly linked to a first inducible promoter sequence;

5 (ii) at least one introduced DNA sequence encoding a protein, polypeptide and/or peptide factor(s) required for growth of the host cell in a protein/serum-free culture medium, expressibly linked to a promoter sequence, the expression of which is regulated by a repressor binding region; and

10 (iii) at least one introduced DNA sequence encoding a repressor molecule which binds to the repressor binding region, expressibly linked to a second inducible promoter sequence, wherein the first and second inducible promoter sequence(s) may be the same or different.

15 18. A host cell according to claim 17, wherein the first and second inducible promoter sequence(s) are the same.

19. A host cell according to claim 17 or 18, wherein the said repressor binding region is a *lac* operator sequence, and said at least one DNA sequence encoding a repressor molecule encodes a *lac* repressor.

20. A host cell according to any one of claims 17 to 19, wherein the first and second inducible promoter sequence(s) is/are selected from the group consisting of the human metallothionein IIA promoter and the modified human metallothionein IIA promoter, M(1)2 and M(2)6.

21. A host cell according to claim 20, wherein the host cell further includes and expresses a DNA sequence encoding a metallothionein.

Sub A1322

A host cell including:

- (i) at least one introduced DNA sequence encoding a protein polypeptide and/or peptide required for growth of the host cell in a protein/serum-free culture medium, expressibly linked to a promoter sequence, the expression of which is regulated by a repressor binding region:

- and
- (ii) at least one introduced DNA sequence encoding a repressor molecule which binds to the repressor binding region, expressibly linked to an inducible promoter sequence.

10

23. A host cell according to claim 22, wherein the said repressor binding region is a *lac* operator sequence, and said at least one DNA sequence encoding a repressor molecule encodes a *lac* repressor.

- 15 24. A host cell according to claim 22 or 23, wherein the inducible promoter sequence(s) is/are selected from the group consisting of the human metallothionein IIA promoter and the modified human metallothionein IIA promoter, M(1)2 and M(2)6.

- 20 25. A host cell according to claim 24, wherein the host cell further includes and expresses a DNA sequence encoding a metallothionein.

Sub A1426

- 25 26. A host cell according to any one of claims 17 to 25, wherein the DNA sequence(s) encoding the protein, polypeptide and/or peptide growth factors(s) encodes a growth factor(s) selected from the group consisting of insulin, modified insulins, insulin-like growth factors, cytokines, mitogenic proteases and mixtures thereof.

T08260-092801

27. A host cell according to claim 26, wherein the DNA sequence(s) encoding the protein, polypeptide and/or peptide growth factor(s) encodes insulin or an insulin-like growth factor.

5 28. A host cell according to claim 26, wherein the DNA sequence(s) encoding the protein, polypeptide and/or peptide growth factor(s) encode insulin or an insulin-like growth factor, and transferrin.

Sub A1 10 29. A host cell according to any one of claims 17 to 28, wherein the mammalian host cell is a Chinese hamster ovary cell.

30. A host cell according to any one of claims 17 to 29, wherein the mammalian host cell is a CHO-K1 cell.

09964338-092801